

9 Meds for Seniors to Avoid

If you or someone you know is over 65, you should be aware of the following drugs that can cause potentially devastating side-effects in older adults. Please print this page out and consult it as needed.

| Opioid Pain Relievers | Heart Medications | Diabetes Drugs |
|---|--|--|
| Demerol, Talwin | Lanoxin +0.125 mg | Diabeta, Micronase, |
| Can cause confusion, | Can be toxic in older adults | Diabinese |
| falls, seizures, confusion and even hallucinations. | and people whose kidneys do not work well. | Can cause severe low |
| | HOL WORK WEIL | blood sugar in older adults. |
| | | |
| NSAIDs | Muscle Relaxants | Estrogen |
| Feldene, Indocin | Flexeril, Robaxin, Soma | Alora, Estrasorb, Cenestrin |
| Can increase the risk of | Can increase the risk of falls, | Can increase your risk |
| indigestion, ulcers & bleeding in the stomach or colon. | and cause constipation, dry mouth & urination problems. | of breast cancer, blood clots & dementia. |
| | | a domonta. |
| | Anti Dovehetie | Anti Anviety and |
| Anticholinergic Drugs | Anti-Psychotic Drugs | Anti-Anxiety and Anti-Insomnia Drugs |
| Elavil, Bentyl, Benadryl | Haldol, Risperdal, Seroquel | Valium, Xanax, Ambien |
| Can cause confusion, constipation, urination problems, blurry vision & low blood pressure. | Can increase the risk of stroke or even death; they can also cause tremors & increase your risk of falls. | Can increase your risk of falls, as well as cause confusion. |



Source: https://geriatricscareonline.org/ProductAbstract/american-geriatrics-society-updated-beers-criteria/CL001